Music in the Brain

T. Christina Zhao
I-LABS, University of Washington
April 2019 at NTU






Music as acoustic signal



- Acoustic features
 - Frequency/Pitch
 - Rhythm/Beat/Meter
 - Spectro-Temporal/Timbre




Today's Topics




- Neural processing of acoustic features in music
- What is the effect of music training?
 - Can it affect music processing?
 - Can the effects generalize to speech?

o Pitch



Timbre




Patil et al, Plos One, 2012



Voice-selective areas

(Belin et al., Nature, 2000)



Rhythm, Beat and Meter





Fig. 1. Illustration of the hierarchy of temporal structure in music.


Zhao et al., Neuropsychologia, 2017



Tracking Isochronous Beat (Geiser et al, J

Neuro, 2012)







Tracking
 Isochronous Beat-- Beta-band Rebound
 (Fujioka et al, J Neuro, 2012)


Processing Meter

(Grahn & Brett, J Cog Neuro, 2007)






Processing Meter – Perceived vs.
 Imaginary (Fujioka et al., J Neuro, 2015)







 Effect on Neural Structure (Schlaug et al., Ann. N.Y Acad. Sci., 2005)




Functional Processing (Pantev et al., Neuro Report,

2001)






o Pitch discrimination using MMR (Terviniemi et al., Euro J Neuro, 2011)




 Pitch discrimination using MMR (Terviniemi et al., Euro J Neuro, 2011)



Meter processing using MMR (Zhao et al,

Neuropsychologia, 2017)





Can it Generalize to Speech?

9-month-old infants



Music Intervention Group (n=22)



Control Group (n=25)


Zhao & Kuhl, PNAS, 2016



Current Study




10-month-olds





Current Study






Current Study



MEG Recordings

Speech Condition







Does the Intervention group exhibit larger MMR


- * to violation in music meter (music condition)?
- * do we observe similar effects in both temporal regions and prefrontal regions between groups?

If so, can the same effects be observed for syllable structure change (speech condition)?

Data Analysis









Data Analysis





Difference wave calculated


Difference wave averaged separately for temporal and prefrontal regions in each individual


















Mixed 2 (between group: exposure vs. control) X 2(within group: temporal vs. prefrontal) ANOVA









Mixed 2 (between group: exposure vs. control) X 2(within group: temporal vs. prefrontal) ANOVA

Discussion



Current study provided strong evidence

- music intervention enhanced temporal structure processing in music
- effect observed in prefrontal regions in addition to temporal regions
- first to show generalization effect to speech domain at 9 month of age



